

Dental Treatment and Breastfeeding Mothers

The information provided is taken from various reference sources. It is provided as a guideline. No responsibility can be taken by the author or the Breastfeeding Network for the way in which the information is used. Clinical decisions remain the responsibility of medical and breastfeeding practitioners. The data presented here is intended to provide some immediate information but cannot replace input from professionals.

There is little research on the safety of dental treatment during breastfeeding but many mothers undertake treatments without harm on a daily basis. The risk of interrupting breastfeeding and substitution of formula appears greater (Dorea 2004).

Fillings

There is no reason to avoid inserting or replacing fillings during breastfeeding. One report suggests that it is prudent to avoid unnecessary removal of fillings during pregnancy or lactation (Barreguard 1995). However there are occasions when a new mother may need a filling inserted or replaced. When mercury is removed some will be vaporised by the high-speed drill and a very small amount may be swallowed or inhaled. These amounts are minute and passage into breastmilk is insignificant compared to the background levels of mercury in the environment. The limitation of the consumption of tuna in line with the FSA guidance is more important on limiting the body burden of mercury and it is very difficult to prove any link between mercury fillings and long term health problems (Lawson). Preventative dental health to minimise the risk of decay is perhaps the message which is of paramount importance.

Local anaesthetic

There is no evidence to interrupt breastfeeding after the use of local anaesthetics. Local anaesthetics work by deadening sensation of the nerve endings around the tooth. There is no evidence of passage into breastmilk and therefore no reason not to continue breastfeeding. The addition of adrenaline to the anaesthetic to reduce bleeding may possibly make a sensitive baby jumpy and irritable so it may be worth asking the dentist to limit use if possible.

Tooth extraction

There is no evidence to interrupt breastfeeding after tooth extraction. If a tooth is to be removed the mother is likely to be offered a local anaesthetic injection or sedation. She may also need pain killers and/or antibiotics.

Sedation/general anaesthetic

There is no evidence to interrupt breastfeeding after sedation or general anaesthesia. The anaesthetics used for dental extractions have a very short half-life (time they act in the body).

By the time the mother is awake most of the drug has been metabolised by the body or exhaled. Similarly sedation with drugs such as midazolam will act for only a short time and by the time the mother is aware of the need to breastfeed once at home, the baby is at worst likely to sleep for a longer than expected period. It may be advisable for another adult to be available to care for the baby until the mother feels totally alert. See fact sheet dental sedation and breastfeeding

To talk to a mum who knows about breastfeeding call the National Breastfeeding Helpline 0300 100 0212

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package.

Analgesics (pain killers)

Breastfeeding mothers may take paracetamol and/or ibuprofen in normal doses after dental treatment.

Antibiotics

Breastfeeding mothers may take antibiotics in normal doses after dental treatment and continue to breastfeed. Babies may have looser bowel motions and may be windy.

Other dental agents

- Mouthwashes – can be used by a breastfeeding mother as they will not be absorbed into the bloodstream e.g. Oraldene®, Corsodyl®, Chlorhexidine mouthwash,
- Gels and liquids for mouth ulcers e.g. Anbesol®, Bonjela®, Adcoryl in Orabase®, Medijel®, Rinstead® pastilles can all be used
- Fluoride toothpastes can be used during breastfeeding. Although there is no research the absorption from the teeth would be minimal and swallowing of excess is unlikely e.g. Durophat®

Tooth whitening

There appears to be no information available on the use of tooth whitening agents during lactation. Whilst it is unlikely that any significant transfer of the agents used into breastmilk will take place, it is unlikely that urgent treatment is necessary and can be delayed until breastfeeding has finished naturally. Unless the products spill from the bath in which the liquid is placed, absorption is unlikely.

References

- Barreguard et al 1995 Occupational and Environmental Medicines
- Dorea JG. Mercury and lead during breast-feeding. Br J Nutr. 2004 Jul;92(1):21-40
- Grandjean et al. Cognitive Deficit in 7 year old children and prenatal mercury exposure 1997 Neurotoxicol Teratol. 1997 Nov-Dec;19(6):417-28 1997
- Peter Lawson BDS, FDS, FRCPS Amalgam and mercury.